

Study of Optokinetic Nystagmus in Meniere's Disease

Davronov U. F., Normuradov N. A., Xo'jamurodov Sh. O.

Department of Otorhinolaryngology No. 2, Samarkand State Medical University

Abstract: The characteristics of optokinetic nystagmus (OKN) were studied in 20 healthy individuals and 30 individuals suffering from Meniere's disease in remission. The results recorded by the method of electronystagmography at frequency optokinetic stimulation 66,90 156 and field/min. found that in healthy people, the frequency of IPOs, the total amplitude and the speed of the slow phase(sp) nystagmus increases proportional to the increase in the frequency optokinetic stimulation; accordingly, consistently reduced the average amplitude window. In Meniere's disease, OKN indicators are generally characterized by lower values than normal, and an increase in the frequency of optokinetic stimulation is accompanied not only by a more pronounced decrease in the total amplitude indicators, but also by a decrease in the speed of the slow phase of OKN. After the glycerol load and with positive results of the glycerol test, the quantitative indicators of OKN increase in patients, approaching the physiological norm. It is concluded that it is appropriate to study the OKN in Meniere's disease to clarify the stage of the disease and predict the effectiveness of treatment.

Key words: Meniere, total amplitude, optokinetic nystagm, glicerol test.

Introduction. According to the literature, the study of optokinetic nystagmus (OKN) is widely used for the differential diagnosis of hemianopsias, oculomotor disorders, and diseases of the Central nervous system [1,2].

In recent years, a number of papers have been published on the study of OKN features in disorders of both Central and peripheral parts of the vestibular analyzer [3,4]. It is experimentally proved that the peripheral labyrinth can give pathological optokinetic nystagmus. Many authors first reported the neutralization of optokinetic nystagmus by supra-threshold post-caloric or post-dictatorial nystagmus in humans and indicated that sub-threshold vestibular stimuli increase the optokinetic response. Visual assessment of OKN disorders under the influence of stimulation in all directions in limes suffering from Meniere's disease and cochlear vestibulopathy caused by vegetative vascular dystonia, cervical osteochondrosis and other causes was not revealed [3,5]. However, when analyzing electronystagmogram was certain violations of the settings window. In this regard, it is advisable to further identify violations of the peripheral parts of the vestibular analyzer, in particular, in Meniere's disease, caused by the hydrops's of the labyrinth.

The aim of the study was to determine the features of the manifestation of optokinetic nystagmus in patients with Meniere's disease.

Materials and Methods: A Total number of examined people in the clinic No. 1 of the Samarkand medical Institute were 50, including 20 healthy (control group) aged 17 to 41 years, who do not suffer from vestibular dysfunction, and 30 people with Meniere's disease aged 25 to 52 years, with a prescription of the disease from 5 months to 14 years. The examination of patients was carried out in the inter-access period. Methods of work inspection, audiometric, vestibulometric study, as well as learning the OCN with the help of stat-tomographic recording. Audiometric examination was performed in a sound-attenuated chamber on an audiometer MA-30 (GDR). hearing tests included tonal threshold, supra-threshold and audiometry.

During vestibulometry, statokinetic stability was determined, and spontaneous and experimental nystagmus obtained with caloric and rotational stimulation was recorded.

OKN was induced by observation of the examinee for optokinetic rotation of the drum, which caused the four black vertical stripes, a Rotating cylinder was carried out at three fixed angular velocities: 99, 135, 2340 in 1 second, which provides the frequency optokinetic stimulation, respectively, 66, 90 and 156 of the bands in the 1 min. In assessing the entry window takes into account the frequency nystagmoid movements, the amplitude and total average speed of nystagmus slow phase (sp) and efficiency bands. In addition, the severity of optosensory and opto vegetative reactions was recorded.

Based on the oral audiometry data, it was found that the examined patients had predominantly unilateral hearing loss on the side of the patient's labyrinth. The curves of the hearing thresholds were mostly horizontal, less often descending and ascending. In most cases, an interval of 10-30 dB was observed between the bone and air conduction curves. We noted low differential thresholds for the perception of sound power (0.4 0.8 dB at an over-threshold tone of 500 and 2000 Hz with an intensity of 20 dB), and a positive SISI test (80-100% of positive responses). According to vestibulometry, the predominant type of disorders was hyporeflexia, in rare cases hyperreflexia of vestibular excitability was observed.

Results. The frequency of OKN in healthy people increased linearly as the frequency of stimulation increased (from 1.2 at the frequency of stimulation of 66 bands/min to 2.64 at the frequency of 156 bands/min). There was an increase in the total amplitude (from 180 to 227.40) and SP (from 18.8 to 26.90/c). At the same time, the average amplitude tended to decrease (from 15 to 8.90) bands (the ratio of the number of nystagmal responses to the number of bands passing in the field of view per unit of time), regardless of the frequency of stimulation, was close to one. Optosensors and opto autonomic reactions were absent (table. 1).

Table 1 characteristics of OKN in normal and Meniere's disease

indicators	Main group			Persons suffering from Meniere's disease		
	Frequency of stimulation(number of bands per 1 minute)					
	66	90	156	66	90	156
Frequency OKN in Hz	1,2±0,1	1,6±0,1	2,64±0,02	1,18±0,01	1,55±0,015	2,43±0,06
The total amplitude (in degrees)	180±8,6	216,6±8	227,4±15,6	123,4±5,8	141,8±6,5	136,2±8,1
The average amplitude (in degrees)	15±0,72	13,7±0,65	8,9±0,55	9,9±0,48	8,3±0,48	5,2±0,29
SMF of optokinetic nystagmus (in degrees / s)	18,8±0,82	23,9±0,88	26,9±0,88	13,1±0,65	15,1±0,76	15,4±0,95
Band efficiency	1±0,1	1±0,1	0,957±0,12	0,952±0,022	0,928±0,016	0,831±0,028

When examining people suffering from Meniere's disease, the WINDOW parameters considered are of a different nature (see table. 1). The frequency of the window increases evenly (from 1.18 to 2.43 Hz). The average amplitude and performance indicators of the bands decrease linearly, and their value is significantly less than the "normal"one. The quantitative values of the total amplitude and CP of nystagmus increase to the frequency of stimulation of 90 bands/min, and with stimulation of 156 bands/min, their indicators decrease slightly, while in the control group, the characteristics under consideration increase linearly. In some patients, an increase in the speed of stimulation caused a feeling of vestibular "discomfort", manifested by dizziness and nausea.

To determine the stage of the process in the subjects, we performed a glycerol test, which is used in diagnostic practice to characterize the reversibility or irreversibility of pathological changes in the receptor formations of the labyrinth in Meniere's disease [5]. The patient was offered to take inside a mixture of pure medical glycerol in half with sour fruit juice at the rate of 1.5 g of glycerol per 1 kg of body weight. Audiometry, vestibulometry, and optokinetic recording were performed up to 3 hours after taking glycerol.

Discussion . According to the results of studies after taking glycerol, patients were divided into two groups. The first group included individuals who had a decrease in tone perception thresholds by 10 to 15 dB during audiometry, and with experimental vestibular samples, the nystagmal response indicators significantly increased. The second group included patients who did not have any significant changes in auditory and vestibular functions after the glycerol test.

Comparison of the results of optokinetic stimulation with the initial data showed that the subjects with a positive glycerol test showed an improvement in all the characteristics of OKN. So, for example, when stimulated 156 hollow/min, the total amplitude increased to 155.5" at the initial value of 136.2", and sp - to 17.8 /s (before taking glycerol, this value was 15.49/s), in patients with a negative glycerol test, we did not note significant changes in the considered characteristics of the OKN.

Comparison of records allows us to conclude that with a positive glycerol test, there is a "revival" of the OKN in comparison with the original data.

Conclusions. Based on the results of the survey, the following conclusion can be made. In healthy people, there is a linear increase in the frequency, total amplitude, and speed of the slow phase of optokinetic nystagmus and a consistent decrease in the average amplitude with an increase in the frequency of optokinetic stimulation. In individuals suffering from Meniere's disease, OKN is characterized by lower parameters, and as the frequency of stimulation increases, the total amplitude of the CMF tends to decrease. Reduced endolymphatic labyrinth as a result of the action is accompanied by an increase in the quantitative characteristics of the WINDOW

Thus, the characteristics of OKN in Meniere's disease may be important for identifying additional criteria for diagnosing diseases of the labyrinth and predicting the effectiveness of treatment.

REFERENCES:

1. Насретдинова М. Т., Карабаев Х. Э. Совершенствование методов диагностики у пациентов с головокружением //Оториноларингология Восточная Европа. – 2017. – Т. 7. – №. 2. – С. 194-198.
2. Nasretdinova M. T., Karabaev H. E., Sharafova I. A. Application of methodologies of diagnostics for patients with dizziness //CENTRAL ASIAN JOURNAL OF MEDICAL AND NATURAL SCIENCES. – 2020. – Т. 1. – №. 1. – С. 29-33.
3. Nasretdinova M. T., Karabaev H. E. Vestibular neuronitis-the problem of systemic dizziness //European science review. – 2019. – Т. 2. – №. 1-2.Singh R.K., Singh M. Otorhinolaryngology Clibics: An International Journal. 2012. Vol. 4(2). P. 81–85.
4. Malik A, Yoshida Y, Erkin T, Salim D, Hamajima N. Hypertension-related knowledge, practice and drug adherence among inpatients of a hospital in samarkand, uzbekistan. Nagoya J Med Sci. 2014;76(3-4):255–263.
5. Wolschner U., Strösser W., Weiser, M., Klein P. Treating vertigo-combination remedy therapeutically equivalent to dimenhydrinate: results of a reference-controlled cohort study. Biol Med. 2010. Vol. 30. P. 184–190.
6. Нормирова, Н. Н., Насретдинова, М. Т., Шадиев, А. Э., & Нормурадов, Н. А. (2023). БОШ АЙЛАНИШИ БИЛАН ОҒРИГАН БЕМОРЛАРДА ПОЗИЦИОН ПАРОКСИЗМАЛ НИСТАГМНИ ЎРГАНИШ. ЖУРНАЛ СТОМАТОЛОГИИ И КРАНИОФАЦИАЛЬНЫХ ИССЛЕДОВАНИЙ, 4(3).

7. Насретдинова, М. Т., Нормирова, Н. Н., Шадиёв, А. Э., & Нормурадов, Н. А. (2023). КОХЛЕОВЕСТИБУЛЯР КАСАЛЛИКЛАРИ БЎЛГАН БЕМОРЛАРДА ВЕСТИБУЛЯР ФУНКЦИЯНИ УРГАНИШ. ЖУРНАЛ СТОМАТОЛОГИИ И КРАНИОФАЦИАЛЬНЫХ ИССЛЕДОВАНИЙ, 4(3).
8. Насретдинова, М. Т., & Нормурадов, Н. А. (2023). Функциональное состояние вестибулярного анализатора по данным вращательной пробы при центральном поражении. Science and Education, 4(8), 65-71.
9. Насретдинова, М. Т., & Нормурадов, Н. А. (2023). Применение консервативной терапии у больных с болезнью Меньера. Science and Education, 4(8), 57-64.
10. Насретдинова, М. Т., & Нормурадов, Н. А. (2023). Оценка диагностических вестибулярных тестов у пациентов с кохлеовестибулярными нарушениями. Science and Education, 4(8), 72-78.
11. Слонова, А. И. (2022). Практики осознанности в развитии творческого потенциала детей: возможности использования в работе практического психолога. Психология. Журнал Высшей школы экономики, 19(3), 642-656.
12. Sayfullayeva Asila Abdulla qizi, R. K. A. qizi ,. (2024). Age-Related Characteristics of Vestibular Neurons. International Journal of Integrative and Modern Medicine, 2(4), 53–56. Retrieved from <https://medicaljournals.eu/index.php/IJIMM/article/view/239>
13. Rasulova, K. (2023). TREATMENT AND PREVENTION OF FUNGAL RHINITIS AND ALLERGIC RHINITIS. Science and innovation, 2(D10), 150-154.
14. Xatamov, J. A., Xayitov, A. A., Boltayev, A. E., & Davronov, U. F. (2023). Comprehensive diagnosis and treatment of chronic purulent otitis media with complications. World Bulletin of Public Health, 28, 73-75.
15. Насретдинова, М. Т., Набиев, О. Р., Хайитов, А. А., & Шадиёв, А. Э. Диагностическое значение нистагма при болезни меньера. междисциплинарный подход по заболеваниям органов головы и шеи, 270.
16. Taxsinovna, N. M., Musinovna, R. K., Farmonqulovich, D. U., & Rahmonovichn, K. I. (2024). VESTIBULAR VASCULAR REACTIONS IN ASSESSMENT OF VESTIBULAR DYSFUNCTION IN PATIENTS WITH CRANIOCEREBRAL TRAUMA. INNOVATIVE ACHIEVEMENTS IN SCIENCE 2024, 3(29), 104-111.
17. Taxsinovna, N. M., Musinovna, R. K., Abruievich, K. J., Maftuna, M., & Ibragimovna, R. E. T. (2024). DIAGNOSTIC INFORMATIVITY OF THE DRUGS USED TO REVEAL INTRALABYRINTHINE HYDROPS ACCORDING TO THE DATA OF AUDIOLOGIC AND BIOCHEMICAL STUDIES. INNOVATIVE ACHIEVEMENTS IN SCIENCE 2024, 3(29), 112-117.
18. Raupova, K., Nasretdinova, M. T., Normuradov, N. A., & Rakhimov, J. H. (2024). TEMPORAL CHARACTERISTICS OF THE ACOUSTIC REFLEXES OF THE INTRA-AURAL MUSCLES IN " NOISE" WORKERS WITH NORMAL HEARING AS WELL AS WITH INITIAL AND PRONOUNCED HEARING IMPAIRMENT. Ethiopian International Journal of Multidisciplinary Research, 11(04), 447-450.